

BRITISH VIEW

MULTIDISCIPLINARY JOURNAL



Anthropologie, Applied Linguistics, Applied Physics, Architecture, Artificial Intelligence, Astronomy, Biological Sciences, Botany, Chemistry, Communication studies, Computer Sciences, Computing technology, Cultural studies, Design, Earth Sciences, Ecology, Education, Electronics, Energy, Engineering Sciences, Environmental Sciences, Ethics, Ethnicity and Racism Studies, Fisheries, Forestry, Gender Studies, Geography, Health Sciences, History, Interdisciplinary Social Sciences, Labour studies, Languages and Linguistics, Law, Library Studies, Life sciences, Literature, Logic, Marine Sciences, Materials Engineering, Mathematics, Media Studies, Medical Sciences, Museum Studies, Music, Nanotechnology, Nuclear Physics, Optics, Philosophy, Physics, Political Science, Psychology, Publishing and editing, Religious Studies, Social Work, Sociology, Space Sciences, Statistics, Transportation, Visual and Performing Arts, Zoology and all other subject areas.

Editorial board

Dr. Marcella Mori Agrochemical Research Centre, Sciensano, Brussels, Belgium.

Dr. Sara Villari Istituto Zooprofilattico Sperimentale della Sicilia, Palermo, Italy.

Dr. Loukia V. Ekateriniadou Hellenic Agricultural Organization, Thessaloniki, Greece.

Dr. Makhkamova Feruza Tashkent Pediatric Medical Institute Uzbekistan

Prof. Dr. Xhelil Koleci Agricultural University of Tirana, Albania.

Prof Dr. Dirk Werling The Royal Veterinary College, London, UK.

Dr. Otabek Yusupov Samarkand State Institute of Foreign Languages

Dr. Alimova Durдона Tashkent Pediatric Medical Institute

Dr. Jamol D. Ergashev Tashkent Pediatric Medical Institute

Dr. Avezov Muhiddin Ikromovich Urgench branch of Tashkent Medical Academy

Dr. Jumaniyozov Khurmatbek Palvannazirovich Urgench state university

Dr. Karimova Aziza Samarkand Institute of Economics and Service

Dr. Rikhsikhodjaeva Gulchekhra Tashkent State Transport University

Dr. David Blane General Practice & Primary Care, University of Glasgow, UK

Dr Raquel Gómez Bravo Research Group Self-Regulation and Health, Institute for Health and Behaviour, Department of Behavioural and Cognitive Sciences, Faculty of Humanities, Education, and Social Sciences, University of Luxembourg, Luxembourg

Dr. Euan Lawson Faculty of Health and Medicine, University of Lancaster, UK

Dr. Krsna Mahbubani General practice, Brondesbury Medical Centre/ University College London, UK

Dr. Patrick Redmond School of Population Health & Environmental Science, King's College London, UK

Dr. Lecturer Liz Sturgiss Department of General Practice, Monash University, Australia

Dr Sathish Thirunavukkarasu Department of Global Health, Population Health Research Institute, McMaster University, Canada

Dr. Sarah White Department of Biomedical Sciences, Macquarie University, New Zealand

Dr. Michael Gordon Whitfield NIHR Health Protection Research Unit in Healthcare-Associated Infections and Antimicrobial Resistance, Imperial College London, UK

Dr. Tursunov Khatam Andijan State Medical Institute Uzbekistan

Manuscripts typed on our article template can be submitted through our website here. Alternatively, authors can send papers as an email attachment to editor@britishview.co.uk

Editor Multidisciplinary Journals

Website: <http://britishview.co.uk>

Email: editor@britishview.co.uk

THE IMPORTANCE OF EARLY REHABILITATION IN THE PROPHYLAXIS OF COMPLICATIONS OF HEMORRHAGIC STROKES

Ruziev Feruz Giyosovich,

Bukhara State Medical Institute

Abstract: Hemorrhagic stroke white survived 75 patients. Psychological, clinical, instrumental and neuropsychological treatment Bilan-birgalikda treatment is the main group of 42 patients of erta rehabilitation, verticalization and hemorrhagic stroke surgery Control and hemorrhagic counterintelligence control and measuring.

Keywords: ischemic stroke, verticalization, early rehabilitation.

INTRODUCTION Hemorrhagic stroke has a poor prognosis, with a mortality rate of 50-70%. The level of damage to brain tissue, the most serious disorders of the main functions of the body are observed in wide blood vessels that affect large parts of the cortex of the cerebral hemispheres. In this case, if a person manages to survive, rehabilitation may take several years, and complete recovery may never occur. BGe and the presence of co-morbidities. The older a person is, the more difficult it is for him to fight a shock, and in the presence of chronic diseases, especially the cardiovascular system, it often becomes impossible. What an attack. Repeated strokes always increase the risk of death. If an elderly person takes an average of 8 months to recover from a first stroke, it can take a year or more to rehabilitate after a second one.

Psychoemotional state of the patient. To fight such a serious disease requires a great will and desire to live, which not every patient has.

It was proved that calcium excretion from the bones increases 3 hours after the patients are immobilized, which in turn causes changes in the bones (B. Izzeky). Nowadays, in the clinics of the western countries, they try to move the patients to a vertical position even when they are in a coma. In the second half of the 20th century, a large number of research studies were conducted on the development of complications such as pneumonia, deep vein thrombosis, and pulmonary artery thrombosis, showing the harmful aspects of bed rest. It has also been found that inadequate reactions to changes in body conditions after several days of strict bed rest are observed, especially in patients with cardiovascular diseases. (Asberg K.H.).

Study objective: Study of the effectiveness of the early rehabilitation process in the prevention of somatic and neurological complications of hemorrhagic strokes.

MATERIALS AND METHODS. Results of examination and analysis of 75 patients who were admitted and treated with the diagnosis of acute cerebral blood circulation disorder, hemorrhagic type in 2022-2023 in the Bukhara branch of the Republican Emergency Medical Scientific Center, emergency neurology and neuroreanimation departments, in order to solve the scientific goals and tasks envisaged by our research work. provided. Patients who underwent early rehabilitation after hemorrhagic stroke Group I (basic group) (BG) consisted of 43 patients, the ratio of women to men was 1:1.2 and the average BGe was 56.3 ± 4.2 , Group II

(comparative) (CG) 32 people, the gender ratio is 1:1.4 with a predominance of women and men, and the average BGe is 58.2 ± 4.9 .

Result and discussion. Subjective, objective, laboratory and instrumental examinations of patients revealed changes that occurred in various degrees in the organs and organ systems. Both groups of patients were given first aid, along with standard medical procedures, verticalization exercises and early rehabilitation measures, including psychological rehabilitation, physiotherapeutic procedures, and therapeutic exercises were recommended. The condition of the patients was evaluated with the help of somatic and neurological complications and quality of life using the NIHSS and Barthel scales on the 1-2 days, 6-7 days, 11-12 days and 17-18 days of the study.

In the main group of patients, parenchymatous hemorrhages accounted for 45.9%, subarachnoid hemorrhages for 32.5%, and subarachnoid-parenchymatous hemorrhages for 21.6%. The results of blood biochemical analysis in research groups are dynamically analyzed and the results before and after the study are presented.

In the research groups, acutely developed neurological deficits are pupil D=S, and anisocoria is not detected, in BG, $n=1$, $1.25 \pm 1.24\%$ of patients have mydriasis.

1- table.

Comparison of pathological changes in the neurological status of patients in research groups.

Indicator	Basic group (n = 43)			Control group (n=32)		
	n	%	m	n	%	m
Diplopia	3	7,0	3,9	1	3,1	3,1
Paresis of the III pair of nerves	1	2,3	2,3	0	0,0	0,0
The face is asymmetrical	18	41,9	7,5	9	28,1	7,9
Dysarthria	21	48,8	7,6	21	65,6	8,4
Aphasia	3	7,0	3,9	3	9,4	5,2
Dysphagia	10	23,3	6,4	5	15,6	6,4
Monoparesis	4	9,3	4,4	3	9,4	5,2
hemiparesis	23	53,5	7,6	21	65,6	8,4
hemiplegia	16	37,2	7,4	8	25,0	7,7

Paresis of oculomotor nerves was observed in $n=1$, $2.3 \pm 2.3\%$ cases in BG, and this condition was not observed in CG. Diplopia BG $n=3$, $7.0 \pm 3.9\%$, CG $n=1$, $3.1 \pm 3.1\%$ is determined. Dysphagia was observed in both groups, BG $n=10$, $23.3 \pm 6.4\%$ and CG $n=5$, $15.6 \pm 6.4\%$, Dysarthria BG $n=21$, $48.8 \pm 7.6\%$, CG $n=21$, $65.6 \pm 8.4\%$, with Aphasia BG $n=3$, $7.0 \pm 3.9\%$ and CG, $n=3$, $9.4 \pm 5.2\%$. Central neuropathy of the facial nerve was observed in BG $n=18$, $41.9 \pm 7.5\%$, CG $n=9$, $28.1 \pm 7.9\%$ cases.

Movement and sensory disorders in the research groups, based on neurological examination data, mild movement disorders were observed in monoparesis type BG

n=3, 3.75±2.12, CG, n=5, 7.14±3.08%, moderate and severe movement disorders hemiparesis BG n=73, 91.25±3.16%, CG n= 62, 88.57±3.8% and hemiplegia BG n=4, 5±2.44%, CG n=3, 4, 29% of cases observed in BG patients with moderate and severe movement disorders can be explained by micro- and macroangiopathic complications of QD. Sensory disorders BG n=69, 86.2± 3.85% are observed in the peripheral part of all muscles. Hemihyperesthesia was observed in BG n=3, 3.75±2.12%, CG n=3, 4.3±2.42% cases.

2- table.**Dynamics of blood biochemical analysis results in research groups.**

Indicator	BG (n=43)	CG (n=32)	p
	M±m	M±m	(p>0,05)
Urea front	7,4±0,45	7,04±0,32	(p>0,01)
Urea last	6,8±0,29	6,18±0,27	(p>0,05)
Creatinine pre	98,1±4,15	97,66±3,65	(p>0,05)
Creatinine last	91,01±2,87	89,79±2,99	(p>0,001)
Glucose pre	6,3±0,39	6,23±0,27	(p>0,01)
Glucose last	8,95±0,32	5,12±0,08	(p>0,05)


Between the main group of patients who underwent early rehabilitation measures and the patients of the comparative group, there were no sharp, significant differences in the results of blood biochemical analysis before and after the study (Table 2). In the practical recommendations as a result of the research, in order to focus the attention of neurologists and neuroreanimatologists on diabetic nephropathy, it was recommended to postpone the verticalization procedure in cases where the creatinine level in the blood is higher than 150 µ.mmol/l and to continue treatment based on the advice of a nephrologist.

Hematomas in the brain were divided into the following groups according to their size:

- Small hemorrhages (hematoma <20ml)
- medium-sized hematoma (hematoma 20-50 ml)
- Large hematoma (hematoma >50 ml)

The frequency of these hematomas in the study groups is reflected in the following diagram in terms of percentages.

Meeting of research groups according to the size of hematomas (%).**1- figure.**



In all patients, hemorrhagic type MSCT of acute blood circulation disorder in the brain is noted. In the course of the study, it was learned that the initiation of a number of early rehabilitation measures directly depends on the laboratory indicators.

2- figure.

Dynamic indicators of complications in the acute period of hemorrhagic stroke





Figure 2 shows the dynamics of complications observed in BG and CG on 1-6 days, 7-12 days and 13-18 days after observation of hemorrhagic stroke. In the first days of the onset of the disease, the symptoms of encephalopathy predominate, and on the 13-18th day of the disease, complications of the respiratory system increase in CG compared to BG, and it is possible to observe the appearance of other complications.

With the NIHSS Scale, the patient's consciousness, vision, motor and sensory function, coordination disorders, gnosis and speech function are evaluated. The NIHSS scale was used to assess the dynamics of the patients' condition in the first and last days of hospitalization and compared between groups.

3-figure.

Dynamics of averBGe indicators on the NIHSS scale.



he study was 5.5 ± 0.29 and CG was 8.2 ± 0.37 before treatment and 4.16 ± 0.29 after rehabilitation measures.

It can be observed that BG was 45.5 ± 2.24 points at the beginning of the study and improved to 69.4 ± 2.05 points at the end of the study. NG, this indicator was 55.5 ± 2.45 at the beginning and 83.7 ± 2.2 at the post-test. It can be observed that the ratio of neurologic deficit level was 1.22 at the beginning of the study and 1.20 at the end of the study.

CONCLUSIONS.

1. To sum up, in the acute period of hemorrhagic stroke, a number of complications are revealed in the somatic and neurological condition of patients and in the dikamika.
2. The results of the early rehabilitation procedures were evaluated using the NIHSS and Barthel scales before and after the study and the improvement of the quality of life was studied.
3. Decrease in developed neurological deficits in the main group and less development of somatic symptoms compared to the comparison group, shows the effectiveness of early rehabilitation measures.

REFERENCES

1. Latysheva V.Ya., Chechetin D.A., Rehabilitation of motor activity of patients in the post-stroke period, // Prakticheskoe rukovodstvo, - Belarus, - Gomel, - 2015, S.15-57.
2. Morozova T.E., Andrushchishina T.B. Arterial hypertension and diabetes mellitus patients - individual selection of antihypertensive medications. Russian Cardiology Journal No. 2 (88) / 2011
3. Ibodullaev Z.R., Stroke and coma.// Tashkent - 2013, S. 17-18. Рузиев, Ф. Г., & Пулатов, С. С. (2020). Совершенство реабилитации больных после инсульта. In Наука и инновации-современные концепции (pp. 90-102).
4. Roziyev F., Khodjaeva N. Cardiovascular system diseases in ischemic strokes BGainst the background of diabetes mellitus //Journal "Meditsina i innovatsii". – 2022. – no. 3. - S. 41-49.
5. Stroke Epidemiology: Advancing Our Understanding of Disease Mechanism and Therapy Bruce OvbiBGele & Mai N. Nguyen-Huynh. The American Society for Experimental NeuroTherapeutics, Inc. 2011.
6. Giyosovich R.F., Sayfullaevich P.S. The score of significant lechebnyx function and early rehabilitation in ischemic stroke and diabetes //British Medical Journal. - 2022. - T. 2. – no. 1.
7. Svetkina A.A. Psychological rehabilitation in hospital with ONMK, // Meditsinskaya psychology v Rossii, — 2016,
8. Wenli Chen, Qian Ye, Xiangtong Ji , Sicong Zhang , Xi Yang. Mirror neuron system based therapy for aphasia rehabilitation, // Frontiers in Psychology, October 2015, Volume 6, P.1-11.
9. Rakhmatova Dilbar Ismatilloeyevna, Narzilloeva Sitora Jakhongirovna The effectiveness of neuroprotective therapy in ischemic stroke // European journal of modern medicine and practice Vol.2 №1, 2022, P. 17-21
10. Рахматова Д.И., Нарзиллоева С.Ж. Диагностика нарушений деятельности центральной нервной системы при ишемическом инсульте с помощью определения когнитивной дисфункции // Тиббиётда янги кун. – Бухара, 2022. - №1(39). - С. 225-229.