

# BRITISH VIEW

MULTIDISCIPLINARY JOURNAL



[www.britishview.co.uk](http://www.britishview.co.uk)

Anthropologie, Applied Linguistics, Applied Physics, Architecture, Artificial Intelligence, Astronomy, Biological Sciences, Botany, Chemistry, Communication studies, Computer Sciences, Computing technology, Cultural studies, Design, Earth Sciences, Ecology, Education, Electronics, Energy, Engineering Sciences, Environmental Sciences, Ethics, Ethnicity and Racism Studies, Fisheries, Forestry, Gender Studies, Geography, Health Sciences, History, Interdisciplinary Social Sciences, Labour studies, Languages and Linguistics, Law, Library Studies, Life sciences, Literature, Logic, Marine Sciences, Materials Engineering, Mathematics, Media Studies, Medical Sciences, Museum Studies, Music, Nanotechnology, Nuclear Physics, Optics, Philosophy, Physics, Political Science, Psychology, Publishing and editing, Religious Studies, Social Work, Sociology, Space Sciences, Statistics, Transportation, Visual and Performing Arts, Zoology and all other subject areas.

### **Editorial board**

**Dr. Marcella Mori** Agrochemical Research Centre, Sciensano, Brussels, Belgium.

**Dr. Sara Villari** Istituto Zooprofilattico Sperimentale della Sicilia, Palermo, Italy.

**Dr. Loukia V. Ekateriniadou** Hellenic Agricultural Organization, Thessaloniki, Greece.

**Dr. Makhkamova Feruza** Tashkent Pediatric Medical Institute Uzbekistan

**Prof. Dr. Xhelil Koleci** Agricultural University of Tirana, Albania.

**Prof Dr. Dirk Werling** The Royal Veterinary College, London, UK.

**Dr. Otabek Yusupov** Samarkand State Institute of Foreign Languages

**Dr. Alimova Durдона** Tashkent Pediatric Medical Institute

**Dr. Jamol D. Ergashev** Tashkent Pediatric Medical Institute

**Dr. Avezov Muhiddin Ikromovich** Urgench branch of Tashkent Medical Academy

**Dr. Jumaniyozov Khurmatbek Palvannazirovich** Urgench state university

**Dr. Karimova Aziza** Samarkand Institute of Economics and Service

**Dr. Rikhsikhodjaeva Gulchekhra** Tashkent State Transport University

**Dr. David Blane** General Practice & Primary Care, University of Glasgow, UK

**Dr Raquel Gómez Bravo** Research Group Self-Regulation and Health, Institute for Health and Behaviour, Department of Behavioural and Cognitive Sciences, Faculty of Humanities, Education, and Social Sciences, University of Luxembourg, Luxembourg

**Dr. Euan Lawson** Faculty of Health and Medicine, University of Lancaster, UK

**Dr. Krsna Mahbubani** General practice, Brondesbury Medical Centre/ University College London, UK

**Dr. Patrick Redmond** School of Population Health & Environmental Science, King's College London, UK

**Dr. Lecturer Liz Sturgiss** Department of General Practice, Monash University, Australia

**Dr Sathish Thirunavukkarasu** Department of Global Health, Population Health Research Institute, McMaster University, Canada

**Dr. Sarah White** Department of Biomedical Sciences, Macquarie University, New Zealand

**Dr. Michael Gordon Whitfield** NIHR Health Protection Research Unit in Healthcare-Associated Infections and Antimicrobial Resistance, Imperial College London, UK

**Dr. Tursunov Khatam Andijan** State Medical Institute Uzbekistan

Manuscripts typed on our article template can be submitted through our website here.

Alternatively, authors can send papers as an email attachment to [editor@britishview.co.uk](mailto:editor@britishview.co.uk)

Editor Multidisciplinary Journals

Website: <http://britishview.co.uk>

Email: [editor@britishview.co.uk](mailto:editor@britishview.co.uk)

## FEATURES OF PREVENTION AND TREATMENT OF SEVERAL CONDITIONS IN CHILDREN OF EARLY AGE.

**Rasulova Nodira Alisherovna**

**Rasulov Alisher Sobirovich**

**Irbutayeva Lola Tashbekovna**

Department of Pediatrics and General Practice, Faculty of Postgraduate Education Samarkand State Medical University

**Abstract:** Rickets is a social problem in pediatrics and is a disease of a growing organism. One of the main causes of seizures in children of the first year of life is a latent vitamin D deficiency. In the city children's hospital No. 1, we examined 65 children who were hospitalized in the intensive care unit for convulsions.

**Keywords:** young children, convulsive conditions, vitamin D, rickets, treatment.

**Relevance.** Seizures are one of the main neurological syndromes of the neonatal period. Their frequency is from 1.1 to 16 per 1000 newborns [3,7]. One of the main causes of seizures in children of the first year of life is a latent vitamin D deficiency and clinically pronounced rickets, in which there is a violation of the formation of cholecalciferol in the skin, a violation of phosphorus-calcium metabolism, a decrease in the formation of active vitamin D metabolites in the liver, kidneys, as well as insufficient intake vitamin D with food [1,4,8]. Rickets is a social problem in pediatrics and is a disease of a growing organism. But in modern conditions, a real deficiency of vitamin D, due to insufficient intake or formation of it in the body, is possible only in early childhood, in most cases in the first year of life, which is objectively related to the nutrition and lifestyle of the child in this age period [2,5,6].

**Purpose of work:** determination of phosphorus-calcium metabolism for the correct appointment of vitamin D in convulsive conditions in children.

**Material and methods:** In the city children's hospital No. 1, we examined 65 children who were hospitalized in the intensive care unit for convulsions. Among them, there were 38 boys and 27 girls. The age of the children ranged from 5 months to 10 months. The children underwent a complete blood count, the level of Ca and P in the blood, as well as EchoEg, and, if necessary, chest X-ray.

**Discussion of the results:** The results of the studies showed that 48 children had a decrease in Ca to 1.7 mmol / l, and in 17 children the level of Ca was on average 2.06 mmol / l. Often the first symptoms of rickets in these children were autonomic disorders. There were such symptoms as sleep disturbance, irritability, tearfulness, excessive sweating.

Due to the presence of an acidotic shift in the body and the acidic nature of sweat, skin irritation has been noted. The most frequent complaints were excessive sweating of the face, especially of the scalp. This, in turn, led to baldness of the back of the head in the majority of children. Urine acquired a sour smell, irritated the child's skin, diaper rash, pyoderma appeared. Compliance of the bones forming the

edges of the large fontanelle was noted in 18 children, curvature of the legs in 13 children, severe muscle hypotension - in 10 children.

From the above, we can conclude that these children did not undergo rickets prophylaxis at the pediatric site.

Upon admission to the hospital, all children were included in the treatment complex vitamin D at a therapeutic dose of 4000 IU per day for 30-45 days. 5 children had repeated convulsions within 2 days. These were children whose blood Ca level was below 1.5 mmol/, and EchoEG showed signs of intracranial hypertension. The inclusion of magnesium sulfate in the treatment complex contributed to the elimination of seizures in this contingent of children. The observations showed that in children who received a sufficient prophylactic dose of vitamin D, even in the presence of intracranial hypertension, convulsions did not recur. All of the above was the basis for revising the methodology for the prevention of rickets. The staff of the Department of Pediatrics developed and implemented a new scheme for the prevention of rickets: Aquadetrim 8 drops or Devaron 6 tablets (4000 IU) 1 time per week from the age of one month to the end of 1 year of life.

Our observations have shown that about 5% of parents have a negative attitude towards the introduction of any drugs to healthy children, about the same number of parents are eager to do “something” to improve the growth and development of the child, and the majority of parents have a positive attitude, but do not realize the importance of the principles prevention and often do not clearly follow the recommendations of the pediatrician. Considering that in our republic there is a system of patronage monitoring of children of the 1st year of life, we recommend that drugs prescribed for preventive purposes be given to the child personally, once a week at the above dose during patronage.

This technique does not increase the workload of nursing staff and, along with it, the doctor is convinced that the child will really receive these drugs.

**Conclusions:** Thus, the problem of convulsive conditions in children caused by rickets does not lose its relevance at the present time. Timely and correctly carried out prevention of rickets is of particular importance. A big mistake is the refusal to prevent rickets with vitamin D, which leads to the development of the disease and, accordingly, to a deterioration in the quality of life of the child: a decrease in motor activity, a violation of the proper formation of the skeletal system, a decrease in immune defense, and a worsening of the course of bronchopulmonary diseases. The scheme developed by us guarantees the obligatory receipt of vitamin D by the child and hence a significant reduction in the convulsive state in children.

In conclusion, it should be emphasized that only complex prevention, including the use of vitamin D, will reduce the incidence of hypocalcemic seizures.

### References

1. Асадуллаев М.М., Расулов Д.К. Клиника, диагностика и особенности течения перинатальных гипоксически-ишемических энцефалопатий // «Неврология». Ташкент, №4 (16).- 2002.- С. 175-176

2. Бадалян Л.О., Темин П.А., Никанорова М.Ю., Медведев М.И. Неонатальная гипокальциемия и гипокальциемические судороги // Педиатрия, 1991.- с.53-58

3. Демин В.Ф. К вопросу о рахите (по поводу статьи Е.В.Неудахина и В.А.Агейкина «Спорные теоретические и практические вопросы рахита у детей на современном этапе») // Педиатрия №4.- 2003.- С. 104-107

4. Коровина Н.А., Захарова И.Н., Чебуркин А.В. Нарушение фосфорно-кальциевого обмена у детей. М., 2005: 70

5. Расулов А.С., Шарипов Р.Х., Дильмурадова К.Р., Мавлянова В.Д., Хатамов Ф.Д., Чальян В.О., Ирбутаева Л.Т. Роль и значение новой программы по профилактике рахита и анемии у детей первого года жизни в оптимизации последипломного образования педиатров //Педиатрия.-Ташкент.-2002.- №2.- с.77-81

6. Романюк Ф.П., Алферов В.П., Колмо Е.А. и др. Рахит (пособие для врачей). — СПб., -2002.- с.61

7. Царегородцева А.В. Современные взгляды на проблему рахита у детей// Педиатрия.М.- 2007.- №6.- с.102-106

8. Шабалов Н.П. Детские болезни. – СПб., 2002. Т.1, С.224-247